

AD-A063 660 MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 7/3
COMPETITIVE RADIATIVE DECAY AND METAL-METAL BOND CLEAVAGE FROM --ETC(U)
JAN 79 J C LUONG, R A FALTYNEK, M S WRIGHTON N00014-75-C-0880
UNCLASSIFIED TR-14 NL

OF
AD
A063680



END
DATE
FILED
3-79
DDC

DDC FILE COPY.

ADA063660

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(2) 1 FVF
REF INSTRUCTIONS
BEFORE COMPLETING FORM

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER ONR-TR-14	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Competitive Radiative Decay and Metal-Metal Bond Cleavage from the Lowest Excited State of Triphenyltin- and Triphenylgermanium-tricarbonyl(1,10-phenanthroline)rhenium		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
7. AUTHOR(s) J. C. Luong, R. A. Faltynek and M. S. Wrighton		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-579
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, Virginia 22217		12. REPORT DATE January 11, 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 11
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited. Reproduction is permitted for any purpose of the United States Government		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Distribution of this document is unlimited.		
18. SUPPLEMENTARY NOTES Prepared for publication in the <u>Journal of the American Chemical Society</u> .		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) optical emission, bond cleavage, quantum yields, lowest excited state, excited state dissociation constant		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <i>(Phi approx. 0.25)</i> The complexes $\text{Ph}_3\text{ERE}(\text{CO})_3(1,10\text{-phenanthroline})$ ($E = \text{Sn, Ge}$) exhibit optical emission (~750 nm) from the lowest excited state in $\text{CH}_2\text{Cl}_2/\text{CCl}_4$ solutions at 298°K and also give E-Re bond cleavage ($\phi = 0.25$) upon photoexcitation under the same conditions. Both the relative emission quantum yield and the reaction quantum yield for $\text{ClRe}(\text{CO})_3(1,10\text{-phenanthroline})$ formation are independent of excitation wavelength consistent with reaction and emission originating from the lowest excited state. The emission is now quenched by CCl_4 in benzene solution, ruling out excited state, bimolecular, electron transfer to CCl_4 as the (over)		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

X
reaction mechanism. Rather, the mechanism is likely slow ($\sim 10^5 \text{ sec}^{-1}$) E-Re dissociation. This notion is in accord with a lowest excited state, which is $(\text{E-Re})\sigma_b \rightarrow (\text{1,10-phenanthroline})\pi^* \text{CT}$ in character.

X (approx. $1/10000 \text{ sec}$)

ACCESSION for

NTIS	White Section
DDC	B. II Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	<input type="checkbox"/>
BY	
DISTRIBUTION/AVAILABILITY COPIES	
D	SPECIAL
A	

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract No. N00014-75-C-0880

9 TECHNICAL REPORT NO. 14

Competitive Radiative Decay and Metal-Metal Bond Cleavage from the
Lowest Excited State of Triphenyltin- and
Triphenylgermanium-tricarbonyl(1,10-phenanthroline)rhenium

by

10 John C. Luong, Robert A. Faltynek, and Mark S. Wrighton

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

11 11 January 1979

12 16 p.

14 TR-14

Reproduction in whole or part is permitted for any purpose of the
United States Government.

Approved for public release; distribution unlimited.

220 007

JOB

Competitive Radiative Decay and Metal-Metal Bond Cleavage from the Lowest
Excited State of Triphenyltin- and Triphenylgermanium-tricarbonyl(1,10-
phenanthroline)rhenium

John C. Luong, Robert A. Faltynek, and Mark S. Wrighton*


Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Abstract: The complexes: $\text{Ph}_3\text{ERe}(\text{CO})_3$ (1,10-phenanthroline) ($\text{E} = \text{Sn, Ge}$) exhibit optical emission (~ 750 nm) from the lowest excited state in $\text{CH}_2\text{Cl}_2/\text{CCl}_4$ solutions at 298°K and also give E-Re bond cleavage ($\phi \approx 0.25$) upon photoexcitation under the same conditions. Both the relative emission quantum yield and the reaction quantum yield for $\text{ClRe}(\text{CO})_3$ (1,10-phenanthroline) formation are independent of excitation wavelength consistent with reaction and emission originating from the lowest excited state. The emission is not quenched by CCl_4 in benzene solution, ruling out excited state, bimolecular, electron transfer to CCl_4 as the reaction mechanism. Rather, the mechanism is likely slow ($\sim 10^5 \text{ sec}^{-1}$) E-Re dissociation. This notion is in accord with a lowest excited state which is $(\text{E-Re})\sigma_b \rightarrow (\text{1,10-phenanthroline})\pi^* \text{CT}$ in character.

Competitive Radiative Decay and Metal-Metal Bond Cleavage from the Lowest Excited State of Triphenyltin- and Triphenylgermanium-tricarbonyl (1,10-phenanthroline)rhenium

Sir:

The rate constants associated with dissociative excited state processes in organometallic complexes have generally not been evaluated, despite the need for such measurements in directly assessing excited state reactivity. There are now a number of reports of bimolecular excited state reaction rate constants for electron and energy transfer.¹⁻⁷ In this communication we report the synthesis and electronic spectral and photochemical characterization of a set of complexes $\text{Ph}_3\text{ERe}(\text{CO})_3\text{L}$ ($\text{E} = \text{Ge}, \text{Sn}$; $\text{L} = 1,10\text{-phenanthroline, 2,2'-bipyridine}$) where it is possible to evaluate $k^*_{\text{dissoc.}}$ associated with the excited state dissociation (1). This represents the first such determination for any metal carbonyl complex, and in particular,



these results bear significantly on the now well-studied metal-metal bonded complexes.⁸⁻¹⁵

Spectral properties for complexes studied are given in the Table; they are prepared by slow addition of a THF solution of Ph_3ECI reaction to a THF solution of the appropriate $[\text{Re}(\text{CO})_3\text{L}]^-$ under Ar. $[\text{Re}(\text{CO})_3\text{L}]^-$ is prepared by the 1% Na/Hg reduction of $\text{ClRe}(\text{CO})_3\text{L}$ in THF solvent. The highly colored E-Re bonded complexes are formed in good yield (~50%) and can be purified by precipitation from concentrated CH_2Cl_2 solution by addition of isoctane. The complexes which have received detailed study to date are for $\text{E} = \text{Ge}$ and Sn and $\text{L} = 1,10\text{-phenanthroline}$ and these two substances have satisfactory

elemental analyses (Alfred Bernhardt, West Germany). Anal. ($C_{33}H_{23}N_2O_3ReSn$); Calcd (Found) % C = 49.52 (49.34); %H = 2.90 (2.92); %N = 3.50 (3.26); ($C_{33}H_{23}N_2O_3ReGe$) %C = 52.54 (52.66); %H = 3.07 (3.12); %N = 3.71 (3.84).

The uv-vis absorption spectra of the complexes Figure 1 and Table are similar to those previously found^{8d} for $(OC)_5Re-Re(CO)_3L$, except for the differences in the energetic position of the bands. We attribute the first absorption band to a metal-metal σ_b to $L\pi^*$ CT transition as in the previously studied Re-Re system. As usual,^{6b,8d,16-18} for such metal carbonyl complexes the CT absorption depends markedly on the solvent; for example the first absorption maximum is at ~440 nm in CH_3CN and ~470 nm in benzene. $Ph_3ERe(CO)_5$ complexes absorb only at wavelengths < 350 nm, ruling out a $\sigma_b \rightarrow \sigma^*$ type transition in the visible; $\sigma_b \rightarrow \sigma^*$ transitions are generally not too solvent sensitive.⁸ The 2,2'-biquinoline complex has been prepared and compared to the others, and the first band is at substantially lower energy in accord with the $(E-Re)\sigma_b \rightarrow L\pi^*$ CT assignment.^{6b,8d}

What is novel is that some of the $Ph_3ERe(CO)_3L$ complexes exhibit detectable emission upon photoexcitation in fluid solution at 25°.

Figure 1. The emission yield is fairly low ($\Phi \sim 10^{-3}$) and the complexes are very photosensitive ($\Phi \approx 0.25$) and the excitation spectrum for wavelengths longer than 300 nm shows that the emission efficiency is wavelength independent. The emission lifetime in degassed 0.5M CCl_4/CH_2Cl_2 solution is 2.6 μ sec and 1.8 μ sec for $Ph_3GeRe(CO)_3$ (1,10-phenanthroline) and $Ph_3SnRe(CO)_3$ (1,10-phenanthroline), respectively, at 25°.

Photoexcitation of the complexes in degassed $\text{CH}_2\text{Cl}_2/\text{CCl}_4$ solutions also results in photoreaction. $\text{ClRe}(\text{CO})_3\text{L}$ is formed quantitatively chemically, and the reaction quantum yields for $\text{Ph}_3\text{GeRe}(\text{CO})_3$ (1,10-phenanthroline) and $\text{Ph}_3\text{SnRe}(\text{CO})_3$ (1,10-phenanthroline) are independent of the excitation wavelength, Table. Thus, it appears that reaction and emission are detectable from the same lowest excited state. Consistently, the emission and reaction are quenched by the triplet quencher anthracene ($E_T = 42 \text{ kcal/mol}$ ¹⁹). The quenching obeys Stern-Volmer kinetics¹⁹ and the Stern-Volmer constant obtained is the same for emission, lifetime, and reaction quenching, Figure 2. This confirms that chemistry and emission result from the same excited state. The fate of the Ge- or Sn-centered fragments has not been quantitatively determined, but E-Cl bonded compounds have been identified as products upon irradiation in the chlorocarbon solutions.

Importantly, it appears that the reaction of the excited state is of the dissociative type and is not one where the excited state reacts bimolecularly with the chlorocarbon. The evidence for this is twofold. First, the complexes emit in benzene solution but concentrations of CCl_4 up to 1M do not quench the emission, while irradiation in the presence of even 0.1M CCl_4 gives clean generation of $\text{ClRe}(\text{CO})_3\text{L}$. Second, irradiation of the reactive complexes in THF and other solvents yields efficient conversion to as yet unidentified products which do not have an E-Re bond as evidenced by bleaching of the low energy visible spectrum. The disappearance quantum yields are similar to those found in the presence of CCl_4 .

Knowing the reaction quantum yield and the emission lifetime under the reaction conditions allows the evaluation of the rate constant (k^*_{dissoc}) for conversion of the excited state to product. For the Ge and Sn complexes studied here the values of $k^*_{\text{dissoc.}}$ are 1.0×10^5 and $1.3 \times 10^5 \text{ sec}^{-1}$, respectively.²⁰ Assuming that the excited state does not relax to a lower state which reacts very rapidly, these can be regarded as being associated with excited state dissociation of the E-Re bond, eq. (1). The excited state rate of E-Re bond cleavage is

very conservatively 10^{11} times the ground state rate constant, since there is no detectable dark reaction over observation times of ~two weeks under conditions where the excited state rate constant is $\sim 10^5 \text{ sec}^{-1}$. Such tremendous enhancement in reaction rate constants is probably common upon excitation of organometallics, but rarely have we had the opportunity to measure the excited state reaction rates. The $(E\text{-Re})\sigma_b \rightarrow L\pi^*\text{CT}$ transition only reduces the σ bond order by one-half, in contrast to the situation in $Mn_2(\text{CO})_{10}$ and related complexes where the $\sigma_b \rightarrow \sigma^*$ can reduce the Mn-Mn bond order to zero.⁸ The $k^*\text{dissoc.}$ values that we have measured are actually quite modest, but it is very likely that the $k^*\text{dissoc.}$ for $\sigma_b \rightarrow \sigma^*$ type excited states would be several orders of magnitude higher than for the $(E\text{-Re})\sigma_b \rightarrow L\pi^*\text{CT}$ excited states studied here.²¹

Acknowledgements.

We thank the National Science Foundation and the Office of Naval Research for support of this research. MSW acknowledges support as a Dreyfus Teacher-Scholar Grant Recipient, 1975-1980. We thank Professors James L. Kinsey and Robert W. Field for valuable discussions and use of the dye laser for lifetime determinations.

John C. Luong, Robert A. Faltynek, and

Mark S. Wrighton*

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

References

1. (a) C. Creutz and N. Sutin, Inorg. Chem., 15, 496 (1976); J. Am. Chem. Soc., 98, 6384 (1976); 99, 241 (1977); (b) N. Sutin and C. Creutz; Adv. Chem. Ser., 168, 1 (1978).
2. (a) V. Balzani, L. Moggi, M. F. Manfrin, F. Bolletta, and G. S. Laurence, Coord. Chem. Rev., 15, 231 (1975); (b) G. S. Laurence and V. Balzani, Inorg. Chem., 13, 2976 (1974); (c) F. Bolletta, M. Maestri, L. Moggi, and V. Balzani, Chem. Comm., 901 (1975); (d) A. Juris, M. T. Gandolfi, M. F. Manfrin, and V. Balzani, J. Am. Chem. Soc., 98, 1047 (1976); (e) R. Ballardini, G. Varani, F. Scandola, V. Balzani, ibid., 98, 7432 (1976).
3. (a) C. R. Bock, T. J. Meyer, and D. G. Whitten, J. Am. Chem. Soc., 96, 4710 (1974); 97, 2909 (1975); (b) R. C. Young, T. J. Meyer, and D. G. Whitten, ibid., 98, 286 (1976); (c) P. J. DeLaise, J. T. Lee, H. Abruna, H. W. Spritschnik, T. J. Meyer, and D. G. Whitten, Adv. Chem. Ser., 168, 28 (1978).
4. (a) H. D. Gafney and A. W. Adamson, J. Am. Chem. Soc., 94, 8238 (1972); (b) J. N. Demas and A. W. Adamson, ibid., 95, 5159 (1973).
5. (a) J. N. Demas, E. W. Harris, C. M. Flynn, Jr., and D. Diemonte, J. Am. Chem. Soc., 97, 3838 (1975); (b) J. N. Demas, D. Diemonte, and E. W. Harris, ibid., 95, 6864 (1973); (c) J. N. Demas, E. W. Harris, and P. P. McBride, ibid., 99, 3547 (1977).
6. (a) M. Wrighton and J. Markham, J. Phys. Chem., 77, 3042 (1973); (b) M. Wrighton and D. L. Morse, J. Am. Chem. Soc., 96, 998 (1974); (c) J. C. Luong, L. Nadjo, and M. S. Wrighton, ibid., 100, 0000 (1978).
7. D. Meisel and M. S. Matheson, J. Am. Chem. Soc., 99, 6577 (1977).
8. (a) H. B. Abrahamson and M. S. Wrighton, Inorg. Chem., 17, 1003 (1978) and J. Am. Chem. Soc., 99, 5510 (1977); (b) M. Wrighton and D. Bredesen, J. Organometal. Chem., 50, C35 (1973); (c) M. S. Wrighton and D. S. Ginley, J. Am. Chem. Soc., 97, 2065, 4246, 4908 (1975); (d) D. L. Morse and M. S. Wrighton, ibid., 98, 3931 (1976).
9. (a) J. L. Hughey, IV., C. R. Bock, and T. J. Meyer, J. Am. Chem. Soc., 97, 4440 (1975); (b) J. L. Hughey, IV, C. P. Anderson, and T. J. Meyer, J. Organometal. Chem., 125, C49 (1977).
10. A. R. Burckett, T. J. Meyer, and D. G. Whitten, J. Organometal. Chem., 67, 67 (1974).
11. R. M. Laine and P. C. Ford, Inorg. Chem., 16, 388 (1977).
12. (a) A. Hudson, M. F. Lappert, P. W. Lednor, and B. K. Nicholson, J. C. S. Chem. Comm., 966 (1974); (b) A. Hudson, M. F. Lappert, and B. K. Nicholson, J. C. S. Dalton, 551 (1977).

13. D. M. Allen, A. Cox, T. J. Kemp, Q. Sultane, and R. B. Pitts, J. C. S. Dalton, 1189 (1976).
14. (a) B. H. Beyers and T. L. Brown, J. Am. Chem. Soc., 97, 947 (1975); 99, 2527 (1977); (b) M. Absi-Halabi and T. L. Brown, ibid., 99, 2982 (1977).
15. C. Giannotti and G. Merle, J. Organometal. Chem., 105, 97 (1976).
16. P. J. Giordano and M. S. Wrighton, Inorg. Chem., 16, 160 (1977), and submitted for publication.
17. H. Saito, J. Fujita, and K. Saito, Bull. Chem. Soc. Japan, 41, 359 and 863 (1968).
18. (a) M. S. Wrighton, H. B. Abrahamson, and D. L. Morse, J. Am. Chem. Soc., 98, 4105 (1976); (b) M. S. Wrighton and D. L. Morse, J. Organometal. Chem., 97, 405 (1975); (c) H. B. Abrahamson and M. S. Wrighton, Inorg. Chem., 18, 0000 (1979).
19. N. J. Turro, "Molecular Photochemistry", Benjamin, New York, 1967.
20. k^* _{dissoc.} = quantum yield for reaction divided by the excited state lifetime. Since there may be geminate recombination of E· and Re·, k^* _{dissoc.} may be regarded as a lower limit but not low by more than a factor of ~4 since the quantum yields are 0.27 and 0.23 for E = Ge and Sn, respectively.
21. Subsequent to the submission of this article an interesting, related paper providing excited state ligand dissociation constants for Rh(III) complexes appeared: M. A. Bergkamp, R. J. Watts, P. C. Ford, J. Brannon, and D. Magde, Chem. Phys. Lett., 59, 125 (1978).

Table - Spectral and Photochemical Properties of $\text{Ph}_3\text{ERe}(\text{CO})_3$ (1,10-phenanthroline) at 25°.

E	Ir Bands, cm^{-1}	(ϵ) ^a	(E-Re) $\sigma_b + \pi^*L$, nm(ϵ) ^b	Em Max, nm (τ , μsec) ^c	$\phi \pm 10\%(\lambda, \text{nm})^d$
Ge	2004 (4870)	450 (4940)	"	765 (2.6)	0.25 (313)
	1900 (3730)				0.30 (366)
					0.27 (436)
					$\frac{0.26(488)}{\text{Av. } = 0.27}$
Sn	2004 (4700)	465 (4610)	"	750 (1.8)	0.26 (313)
	1903 (3840)				0.25 (366)
					0.22 (436)
					$\frac{0.20(488)}{\text{Av. } = 0.23}$

^a In CH_2Cl_2 with 0.5 M CCl_4 at 298°K measured with a PE 180 with matched path cells.

^b In CH_2Cl_2 with 0.5 M CCl_4 at 298°K measured using a Cary 17 uv-vis-nir spectrophotometer.


^c Corrected emission maximum in CH_2Cl_2 with 0.5M CCl_4 using a PE Model MPF44 exciting at 480 nm. The lifetime was measured using either a rhodamine dye laser (560 nm, ~5 nsec pulse width) or a Xenon Corporation Model 437 Nanopulser excitation source filtered to pass 436 nm light and the detection optics of a TRW Decay Time Fluorometer with output to an oscilloscope.


^d Reaction quantum yield for disappearance of E-Re species is equal to the formation quantum yield for $\text{CIRe}(\text{CO})_3$ (1,10-phenanthroline) which exhibits ir absorptions (ϵ) in $\text{CH}_2\text{Cl}_2/0.5\text{M}$ CCl_4 at 2024 (4600), 1922 (3015) and 1898 (2600). All ϕ 's are in degassed $\text{CH}_2\text{Cl}_2/0.5\text{M}$ CCl_4 solutions at $\sim 2 \times 10^{-3}$ M E-Re species. Light intensities are $\sim 3 \times 10^{-7}$ ein/min.

Figure 1. Absorption and emission of $\text{Ph}_3\text{GeRe}(\text{CO})_3$ (1,10-phenanthroline).

The emission spectra are corrected for variation in detector sensitivity and are excited at 480 nm. The sensitivity for the 298 and 77°K emission is not the same, and the absorption spectral change from 298 to 77°K is not corrected for solvent contraction.

Figure 2. Stern-Volmer plots for anthracene quenching of emission intensity (\circ), emission lifetime (\bullet) and reaction quantum yield (Δ) for 2×10^{-3} M $\text{Ph}_3\text{GeRe}(\text{CO})_3$ (1,10-phenanthroline) at 298°K in a degassed CH_2Cl_2 solution of 0.5 M CCl_4 .





For NR 051-579

472:GAN:716:tam
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314 12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP 1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney 1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division 1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko 1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940 1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380 1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller 1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division 1
	1	Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division 1

Encl

TECHNICAL REPORT DISTRIBUTION LIST, 051A

<u>No.</u>	<u>Copies</u>	<u>No.</u>	<u>Copies</u>
Dr. M. A. El-Sayed University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1	Dr. M. Rauhut American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805	1
Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163	1	Dr. J. I. Zink University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1
Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. B. Schechtman IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. J. R. MacDonald Naval Research Laboratory Chemistry Division Code 6110 Washington, D.C. 20375	1		
Dr. G. B. Schuster University of Illinois Chemistry Department Urbana, Illinois 61801	1		
Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1		
Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007	1		
Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1		